

DEPARTMENT OF HEALTH & HUMAN SERVICES

Public Health Service

National Institutes of Health
Bethesda, Maryland 20892
www.nih.gov

OCT 07 2004

The Honorable Henry A. Waxman
Ranking Minority Member
Committee on Government Reform
House of Representatives
Washington, D.C. 20515-6143


Dear Mr. Waxman:

Thank you for your letter, co-signed by Representative Elijah E. Cummings, in which you express concern regarding the presentation of scientific evidence on the efficacy of syringe exchange and "harm reduction" programs to prevent the spread of HIV and other blood borne illnesses. I have enclosed a brief response to the areas of interest you identified in your letter.

I hope you find this information useful. If you have further questions or need additional information, please contact Dr. Steve Gust, Interim HIV/AIDS Coordinator at the National Institute on Drug Abuse, at 301-443-6480.

An identical letter is being sent to Mr. Cummings.

Sincerely,



Elias A. Zerhouni, M.D.
Director

Enclosure

17544

1. Strategies That Have Proven Successful in Reducing The Risk of HIV Infection Among IDUs.

One successful strategy for reducing the risk of HIV among injection drug users (IDUs) is to provide drug abuse treatment. Drug treatment programs provide a good setting for reaching IDUs and their partners with HIV prevention and care messages and interventions. It also can be a bridge to other needed services, such as primary health care, mental health, or other social services.

Numerous studies, primarily focused on methadone maintenance treatment (MMT), have shown that substance abuse treatment programs can have a dramatic effect on HIV transmission among opiate injectors, reducing their risk as much as 4- to 6-fold.^{1,2} Drug abuse treatment works principally because it helps IDUs decrease the number of injections or helps them stop injecting altogether. Less use leads to fewer drug-related risk behaviors, and that in turn leads to fewer exposures to HIV. The beneficial effects of MMT are most evident when treatment lasts a sufficiently long time and when methadone doses are high enough to effectively block drug craving: One study showed that 3.5 percent of methadone patients who had been in treatment continuously for 18 months had become infected with HIV, compared to 22 percent of out-of-treatment IDUs;³ another study showed that at 36 months, 8 percent of IDUs in treatment had become infected,⁴ as compared to 30 percent of injectors not in treatment. An analysis of 20 years of social and medical data on 622 MMT patients in New York City showed that those patients who received methadone doses of 80 mg or more were significantly less likely to have HIV infection than patients who received smaller doses.⁵ The protective value of higher doses was independent of a number of other risk factors, including year of last cocaine injection, needle sharing in shooting galleries, number of IDU sex partners, income, and race/ethnicity. Moreover, among non-injection cocaine users, drug treatment has also been shown to decrease cocaine use from an average of 10 days per month at baseline to 1 day per month at 6 months. Reduction in cocaine use was associated with an average 40 percent decrease in HIV risk across gender, and ethnic groups, mainly as a result of fewer sexual partners and less unprotected sex.⁶

Drug addiction treatment⁷ is an essential component of a comprehensive prevention program to reduce risk of HIV and other blood-borne infections among IDUs. Since the late 1980s, studies have shown that treatment works because drug users in treatment stop or reduce their drug use and related risk behaviors, including use of non-sterile syringes and unsafe sex. Drug treatment programs also serve an important role in providing up-to-date information on HIV/AIDS, hepatitis, and other sexually transmitted diseases (STDs), counseling and testing services for these infections, and referrals for their clients to obtain medical and social services.

However, the majority of those needing treatment are not currently in a treatment program. The NIDA Community-Based Outreach Model^{8,9} was designed to reach out-of-treatment IDUs who are unable or unwilling to stop using and injecting drugs and who cannot or will not access drug treatment. Compared to those in treatment, out-of-treatment IDUs are at significantly greater risk of HIV and other infections because they are more likely to inject drugs more frequently, to share drugs, syringes, and other injection equipment, and to practice unsafe sex while under the influence of drugs. The outreach program developed by NIDA attempts to reduce HIV risk through education on the risk factors for HIV transmission and by teaching effective skills in reducing those risks.

The Federal Government has extensively examined the effectiveness of syringe exchange programs (SEPs) dating back to 1993, including reviews by the Government Accountability Office.¹⁰ Several non-governmental organizations, including the American Psychiatric Association, and others have also endorsed the use of SEPs as effective public health interventions. The current scientific literature supports the conclusion that SEPs can be an effective component of a comprehensive community-based HIV prevention effort.

References:

- ¹Joseph H, Stancliff S, Langrod J. Methadone maintenance treatment (MMT): a review of historical and clinical issues. *The Mt. Sinai Journal of Medicine* 67(5&6):347-364, 2000.
- ²Metzger DS, Navaline H, Woody GE. Drug abuse treatment as AIDS prevention. *Public Health Reports* 113(Suppl 1):97-106, 1998.
- ³Metzger DS, Woody GE, McLellan AT, O'Brien CP, Druley P, Navaline H, De Philippis D, Stolley P, Abrutyn E. Human immunodeficiency virus seroconversion among intravenous drug users in- and out-of-treatment: An 18-month prospective follow-up. *Journal of Acquired Immune Deficiency Syndromes*, 6, 1049-1056, 1993.
- ⁴McLellan AT, Metzger DS, Alterman AI, Woody GE, Durell J, O'Brien CP. Evaluating the effectiveness of addiction treatment: Reasonable expectations, appropriate comparisons. *Milbank Quarterly*, 74, 51-85, 1996.
- ⁵Hartel DM, Schoenbaum EE. Methadone treatment protects against HIV infection: two decades of experience in the Bronx, New York City. *Public Health Reports* 113(Suppl 1):107-115, 1998.
- ⁶Woody GE, Gallop R, Luborsky L, Blaine J, Frank A, Salloum IM, Gastfriend D, Crits-Christoph P. HIV risk reduction in the National Institute on Drug Abuse Cocaine Collaborative Treatment Study. *J Acquir Immune Defic Syndr* 33(1):82-87, 2003.
- ⁷NIDA. Principles of Drug Addiction Treatment: A Research-Based Guide. NIH Publication No. 99-4180, Oct 1999.
- ⁸NIDA. Principles of HIV Prevention in Drug-Using Populations: A Research-Based Guide. U.S. Department of Health and Human Services, National Institutes of Health, National Institute on Drug Abuse; NIH Publication No. 02-4733, 2002.
- ⁹NIDA. The NIDA Community-Based Outreach Model: A Manual to Reduce the Risk of HIV and Other Blood-borne Infections in Drug Users. U.S. Department of Health and Human Services, National Institutes of Health, National Institute on Drug Abuse; NIH Publication No. 00-4812, 2000.
- ¹⁰U.S. General Accounting Office. Needle Exchange Programs: Research Suggests Promise as an AIDS Prevention Strategy. Report No. GAO/HRD-93-60. Washington, DC: US GPO, 1993.

2. The Role Played by Harm Reduction Programs in Stemming the Spread of HIV in the United States.

3. The Relative Rate of HIV Infection in Cities That Have Implemented Harm Reduction Programs Versus Those That Have Not.

As a public health agency, the goal of the National Institutes of Health (NIH) and specifically the National Institute on Drug Abuse (NIDA) is to improve the quality of the Nation's addiction treatment and prevention, using science as the vehicle. The term 'harm reduction' has various meanings depending upon the context in which it is used, and is not viewed as a scientific term for any particular approach to addressing drug addiction. However, a great deal of research has been conducted on methods of reducing risks to health, such as syringe exchange programs (SEPs).

Research shows that SEPs, when implemented as part of a comprehensive HIV/AIDS prevention strategy, can be an effective public health approach to reduce the spread of HIV and other blood borne pathogens in the community. SEPs reduce the circulation time of contaminated injection equipment and thereby reduce opportunities for reuse of contaminated injection equipment and the transmission of new infections.¹⁻² A number of studies conducted in the U.S. have shown that SEPs do not increase drug use among participants or surrounding community members and are associated with reductions in the incidence of HIV, hepatitis B, and hepatitis C in the drug-using population.³⁻⁷

Hurley, et al.,⁸ reviewed published and unpublished reports from 1984 to 1994 on HIV seroprevalence among IDUs in 81 cities across Europe, Asia, and North America with and without SEPs. On average, seroprevalence increased by 5.9 percent per year in the 52 cities without SEPs and decreased by 5.8 percent per year in the 29 cities with SEPs. The average annual change in seroprevalence was 11 percent lower in cities with SEPs. Thus, in cities with SEPs, HIV seroprevalence among IDUs decreased on average, but in cities without SEPs, HIV seroprevalence increased, suggesting that SEPs led to a reduction in HIV incidence among IDUs.

References:

- ¹Kaplan, E.H., & Heimer, R. HIV incidence among needle exchange participants: Estimates from syringe tracking and testing data. *J AIDS & Human Retrovirol*, 7(2):182-189, 1994.
- ²Kaplan, E.H. Operational modeling of needle exchange programs. In: *Proceedings, Workshop on Needle Exchange and Bleach Distribution Programs*: 202-249. National Research Council and Institute of Medicine. Washington, DC: National Academy Press, 1994.
- ³Kaplan, E.H., Khoshnood K., and Heimer R. A decline in HIV-infected needles returned to New Haven's needle exchange program: client shift or needle exchange? *Am J Public Health*, 84(12): 1991-1994, 1994.
- ⁴Des Jarlais, D.C., Marmor, M., Paone, D., Titus, S., Shi, Q., Perlis, T., Jose, B., & Friedman, S.R. HIV incidence among injecting drug users in New York City syringe-exchange programmes. *The Lancet*, 348:987-991, 1996.
- ⁵Hagan, H., Des Jarlais, D.C., Friedman, S.R., Purchase, D., & Alter, M.J. Reduced risk of hepatitis B and hepatitis C among injecting drug users participating in the Tacoma syringe exchange program. *Amer J Public Health*, 85(11):1531-1537, 1995.

- ⁶Kaplan, E.H., & Heimer, R. HIV incidence among New Haven needle exchange participants: Updated estimates from syringe tracking and testing data. *J AIDS & Human Retrovirol*, 10(2):175-176, 1995.
- ⁷Vlahov, D., Junge, B., Brookmeyer, R., Cohn, S., Riley, E., Armenian, H., & Beilenson, P. Reductions in high risk drug use behaviors among participants in the Baltimore needle exchange. *J AIDS & Human Retrovirol*, 16(5):400-406, 1997.
- ⁸Hurley, S., Jolley, D.J., Kaldor, J.M. Effectiveness of needle-exchange programmes for prevention of HIV infection. *The Lancet* 349:1797-1800, 1997.

4. Evidence Comparing HIV Treatment Regimen Compliance Among IV Drug Users vs. Non-IV Drug Users.

HIV-infected drug abusers can achieve positive health outcomes if they have access to and adhere to treatment with antiretroviral drugs (ART). Studies have also demonstrated the importance of ongoing interventions to reduce drug abuse and associated risk behaviors in order to maximize the health benefits of ART. However, there is cause for concern that health outcomes in drug abusers infected with HIV may be inferior to non-drug users. The often chaotic lifestyles of drug abusers combined with their increased likelihood of co-occurring medical and psychiatric conditions can complicate their treatment and prevent their achieving the same health outcomes as non-drug users. Access to medical care is another crucial factor. Individuals who receive HIV treatment later in the course of their disease are more likely to have viral rebound associated with development of resistance to ART than those who receive early treatment.¹ Finally, preclinical or basic research studies indicate that some drugs of abuse affect the immune system, the target of HIV infection, which may also impact vulnerability to infection and course of illness.

Factors associated with treatment compliance in drug abusing populations are discussed below.

Adherence to HIV treatment among drug users

The cumulative research indicates that non-adherence to antiretroviral therapy (ART) occurs in both drug users and non-drug users, reflecting the difficulty of adhering to complex regimens which require high accuracy in dosing schedule and compliance with dietary instructions. Estimates are that about 40 percent of patients receiving ART have significant problems with adherence.² A study of adherence among non-drug-using patients found 53.1% reported taking all medication on time according to dietary instructions, i.e., were fully adherent.³ It is important to recognize that not only do treatment outcomes depend upon adherence to medication regimens, but also the risk of developing resistant HIV strains may be related to the level of sustained treatment adherence.¹

A number of predictors of poor ART treatment adherence have been demonstrated in research studies. These include illicit drug use, as well as depression, alcohol use, poor self-efficacy, and certain health beliefs. However, the evidence from individual studies is not consistent--in some cases no differences are found between drug users, former drug users and non-drug users, and in other cases clear evidence of poorer adherence and lower HIV viral suppression is found in active drug users. Examples of this research follow:

- In one study, the strongest predictor of poor ART adherence in drug users was active cocaine use (27% in abstinent users vs. 68% in active users). Other factors included female gender, being unmarried, screening positive for depression and use of alcohol.⁴
- In a cohort of HIV infected women adherence was found not to be stable over time, with factors such as active drug/alcohol use, more frequent antiretroviral dosing, younger age, and lower initial CD4 lymphocyte count predicting poor ART adherence.⁵
- Lucas, et al.,⁶ identified the effects of substance abuse status on utilization of highly active anti-retroviral therapy (HAART), medication adherence, and virologic and immunologic responses to therapy in a cohort of HIV-1-infected patients attending an urban HIV clinic. Active drug use was strongly associated with underutilization of HAART, non-adherence,

and inferior virologic and immunologic responses to therapy. Former drug users and non-drug users were similar in all outcomes.

- Another study by this group⁷ indicated that switching from non-use to substance abuse was strongly associated with worsening ART use and adherence, and less frequent HIV-1 RNA suppression, compared to remaining free of substance abuse. Conversely, switching from substance abuse to non-use was strongly associated with improvements in ART use, adherence and treatment outcomes.
- Not all studies support an association between drug abusers and poor adherence. A study of factors relating to adherence to antiretroviral therapy among pregnant women indicated that adherence to antiretroviral therapy was not significantly associated with use of illicit drugs. Analyses were based on pharmacy claims data in a sample of 549 HIV-infected women who were prescribed antiretroviral therapy and who delivered live infants.⁸

References:

- ¹Sethi AK, Celentano DD, Gange SJ, Moore RD, Gallant JE. Association between adherence to antiretroviral therapy and human immunodeficiency virus drug resistance. *CID* 37:1112-1118, 2003.
- ²Cheever LW, Wu AW. Medication adherence among HIV-infected patients: understanding the complex behavior of patients taking this complex therapy. *Curr Infect Dis Rep* 4:401-407, 1999.
- ³Nieuwkerk PT, Sprangers AG, Burger DM et al, Limited patient adherence to highly active antiretroviral therapy for HIV-1 infection in an observational cohort study. *Arch Intern Med* 161:1962-1968, 2001.
- ⁴Arnstén JH, Demas PA Grant RW, Gourevitch MN, FarzadeganH, Howard AA, Schoenbaum EE. Impact of active drug use on antiretroviral therapy adherence and viral suppression in HIV-infected drug users. *J Gen Intern Med* 17(5):377-381, 2002.
- ⁵Howard AA, Arnsten JH, Yungtai L, Vlahov D, Rich JD, Schuman P, Stone VE, Smith DK, Schoenbaum EE. A prospective study of adherence and viral load in a large multi-center cohort of HIV-infected women. *AIDS* 16:2175-2182, 2002.
- ⁶Lucas GM, Cheever LW, Chaisson RE, Moore RD. Detrimental effects of continued illicit drug use on the treatment of HIV-1 infection. *J Acquir Immune Defic Syndr* 27:251-259, 2001.
- ⁷Lucas GM, Gebo KA, Chaisson RE, Moore RD. Longitudinal assessment of the effects of drug and alcohol abuse on HIV-1 treatment outcomes in an urban clinic. *AIDS* 16:767-774, 2002.
- ⁸Laine C, Newschaffer CJ, Zhang D, Cosler L, Hauck WW, Turner BJ. Adherence to antiretroviral therapy by pregnant women infected with human immunodeficiency virus: a pharmacy claims-based analysis. *Obstetrics and Gynecology* 95:167-173, 2000.

5. The Use of Harm Reduction Strategies in Areas Other Than HIV and Drugs, Such as Speed Limits, Seat-Belt Laws, Minimum Age of Alcohol Consumption, and Public Education and Peer Outreach Concerning Smoking.

The reduction of risk for injury and death has been the focus of research in a number of fields. In traffic safety, reduced speed limits¹ and seat-belt laws² have reduced the likelihood of crashes and the severity of injuries sustained in those crashes. The Insurance Institute for Highway Safety has published a selection of findings on the prevention and consequences of increased speed limits in several editions of *Status Report* including, "Seven straight years: deaths higher after 65 mph speed limits than before" in 1994 and "Faster travel and the price we pay" in 2003. More information is available on the Insurance Institute for Highway Safety's website at <http://www.highwaysafety.org>. The Centers for Disease Control and Prevention's (CDC) National Center for Injury Prevention and Control houses a Task Force on Community Preventive Services which has published findings on seat-belt use interventions and the effectiveness of safety belt use laws. CDC's reports have been featured in publications including numerous issues of the *Morbidity and Mortality Weekly Report* (MMWR) and Volume 21 of the *American Journal of Preventive Medicine* (AJPM).³ More information is available on the National Center for Injury Prevention and Control's website at <http://www.cdc.gov/ncipc/>.

Research in the alcohol field has shown that crashes and injuries have been reduced by raising the drinking age,^{4,5} reducing the allowable blood alcohol concentration (BAC) for drivers,⁶ and enacting zero tolerance laws for younger drivers.^{7,8} Research has also shown that providing a brief intervention to reduce a person's drinking lowers the probability of making a subsequent visit to an emergency room.⁹

Education aimed at better informing the public on smoking and health issues are an important part of tobacco control and prevention efforts.¹⁰ It is vital that the public understand that, to date, the only proven way to reduce the enormous burden of disease and death due to tobacco use is to prevent youth from beginning to smoke, and to help smokers, both youth and adults to quit.¹¹ Today, we have much to offer people who smoke and want to quit, including effective behavioral treatments and medications.¹² The evidence strongly suggests that people who keep trying to quit do succeed, although many will require numerous attempts before being successful.¹³

Recently, a number of new tobacco products with claims purported to reduce health risk have entered the market.¹⁴⁻¹⁶ Unlike smoking cessation products, tobacco products do not undergo rigorous, objective scrutiny either for their constituents or for the accuracy of their health claims. A greater science base is required before we will know what effect these new products will have on the health of the public.¹⁷⁻¹⁹

To be effective, education, and outreach efforts must take into account the knowledge, attitudes, and behaviors – among other factors – of the intended audience.²⁰ To understand these and related issues, the National Cancer Institute (NCI) has developed and implemented the Health Information National Trends Survey (HINTS), which collects nationally representative data about the American public's use of cancer-related information and perception of cancer risks. HINTS contains questions about tobacco product use, including tobacco products purported to reduce health risk. These data will be useful to help shape future public education efforts.²¹

References:

- ¹Transportation Research Board, National Research Council, Committee for Guidance on Setting and Enforcing Speed Limits. *Managing Speed: Review of Current Practice for Setting and Enforcing Speed Limits*. Special Report #254, National Academy Press, Washington, D.C., 1998.
- ²Shults RA, Elder RW, Sleet DA, Thompson RS, and Nichols JL. Primary enforcement seat belt laws are effective even in the face of rising belt use rates. *Accident Analysis and Prevention* 36:491-493, 2004.
- ³"Reviews of Evidence Regarding Interventions to Increase the Use of Safety Belts". *American Journal of Preventive Medicine* 2001;21:4S:48-65.
- ⁴Shults RA, Elder RW, Sleet DA, Nichols JL, Alao MO, Carande-Kulis VG, Zaza S, Sosin DM, Thompson RS. Reviews of evidence regarding interventions to reduce alcohol-impaired driving. *American Journal of Preventive Medicine* 21:66-88, 2001.
- ⁵Wagenaar AC, Toomey TL. Effects of minimum drinking age laws: Review and analysis of the literature from 1960 to 2000. *Journal of Studies on Alcohol*. 14(Supplement): 206-225, 2002.
- ⁶Jonah B, Mann R, Macdonald S. The effects of lowering legal blood alcohol limits: A review. In: *Proceedings of the 15th International Conference on Alcohol, Drugs and Traffic Safety*. Stockholm, Sweden, 2001.
- ⁷Hingson R, Heeren T, Winter M. Lower legal blood alcohol limits for young drivers. *Public Health Reports* 109:738-744, 1994.
- ⁸Wagenaar AC, O'Malley PM, LaFond C Lowered legal blood alcohol limits for young drivers: Effects on drinking, driving and driving after drinking behaviors in 30 states. *American Journal of Public Health* 91:801-804, 2001.
- ⁹Gentilello LM, Rivara FP, Donovan DM, Jurkovich GJ, Daranciang E, Dunn CW, Villaveces A, Copass M, Ries RR. Alcohol interventions in a trauma center as a means of reducing the risk of injury recurrence. *Annals of Surgery* 20:473-483, 1999.
- ¹⁰Centers for Disease Control and Prevention. *Best Practices for Comprehensive Tobacco Control Programs*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, August 1999.
- ¹¹U.S. Department of Health and Human Services. *Reducing Tobacco Use: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2000.
- ¹²Fiore MC, Bailey WC, Cohen SJ, et al. *Treating Tobacco Use and Dependence*. Clinical Practice Guideline. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, June 2000.
- ¹³Centers for Disease Control and Prevention. Cigarette smoking among adults – United States, 1998. *Morb Mortal Wkly Rep*. 2000;49:881-884.
- ¹⁴Buchhalter, A.R. and Eissenberg, T. Preliminary evaluation of a novel smoking system: Effects on subjective and physiological measures and on smoking behavior. *Nicotine & Tobacco Research* 2: 39-43, 2000.
- ¹⁵Breland, A.B., Buchhalter, A.R., Evans, S.E., and Eissenberg, T. Evaluating acute effects of potential, reduced-exposure products for smokers: Clinical laboratory methodology. *Nicotine & Tobacco Research*, S131-S140, 2002.

- ¹⁶Breland, A.B., Acosta, M.C., and Eissenberg, T. Tobacco-specific nitrosamines and potential reduced exposure products for smokers: A preliminary evaluation of AdvanceTM. *Tobacco Control*, 2003.
- ¹⁷Stratton K.S., Wallace R., Bondurant S., (eds.). *Clearing the Smoke: Assessing the Science Base for Tobacco Harm Reduction*. Washington, DC: National Academy Press, 2001.
- ¹⁸Hatsukami, D.K., Slade, J., Benowitz, N.L., Giovino, G., Gritz, E.R., Leischow, S., and Warner, K. E. Reducing tobacco harm: research challenges and issues. *Nicotine & Tobacco Research*, S73-S85, 2002.
- ¹⁹Fowles, J., and Dybing, E. Application of toxicological risk assessment principles to the chemical constituents of cigarette smoke. *Tobacco Control* 12: 424-430, 2003.
- ²⁰U.S. Department of Health & Human Services. Making Health Communication Programs Work. Public Health Service, National Cancer Institute. NIH Publication No. 02-5145. Revised December 2001. Printed September 2002 (Page 219).
- ²¹Health Information National Trends Survey. <http://dccps.nci.nih.gov/hints/index.html>.

September 2004